

# Measuring the threat from a distance: insight into the complexity and perspectives for implementing sentinel plantation to test the host range of *Xylella fastidiosa*

Noemi Casarin<sup>1</sup>, Séverine Hasbroucq<sup>2</sup>, Júlia López-Mercadal<sup>3</sup>,  
Miguel Ángel Miranda<sup>3</sup>, Claude Bragard<sup>1</sup>, Jean-Claude Grégoire<sup>2</sup>

**1** Earth and Life Institute Applied Microbiology (ELIM), Université Catholique de Louvain (UCLouvain), Croix du Sud 2 bte L7.05.03, 1348 Louvain-la-Neuve, Belgium **2** Spatial Epidemiology lab (SpELL), Université Libre de Bruxelles (ULB), CP 160/12, 50 av. F.D. Roosevelt, 1050 Bruxelles, Belgium **3** Zoologia Aplicada i de la Conservació (ZAP), Universitat de les Illes Balears (UIB), Cra. De Valldemossa, km 7.5, 07122 Palma de Mallorca, Illes Balears, Spain

Corresponding authors: Jean-Claude Grégoire ([jean-claude.gregoire@ulb.be](mailto:jean-claude.gregoire@ulb.be)), Claude Bragard ([claude.bragard@uclouvain.be](mailto:claude.bragard@uclouvain.be))

Academic editor: Alberto Santini | Received 6 July 2022 | Accepted 8 November 2022 | Published 18 May 2023

**Citation:** Casarin N, Hasbroucq S, López-Mercadal J, Miranda MÁ, Bragard C, Grégoire J-C (2023) Measuring the threat from a distance: insight into the complexity and perspectives for implementing sentinel plantation to test the host range of *Xylella fastidiosa*. In: Jactel H, Orazio C, Robinet C, Douma JC, Santini A, Battisti A, Branco M, Seehausen L, Kenis M (Eds) Conceptual and technical innovations to better manage invasions of alien pests and pathogens in forests. NeoBiota 84: 47–80. <https://doi.org/10.3897/neobiota.84.90024>

## Abstract

The sentinel plantation concept consists of assessing the impact of exotic factors, such as pests and pathogens, on plants of interest by planting them out of their native range. This tool is a way to enhance knowledge for pest risk analysis (PRA) by guiding decisions on how quarantine organisms should be regulated and where to focus prevention and surveillance efforts for an early detection. In this study, the sentinel method was used in the case of research on *Xylella fastidiosa*, a plant pathogenic bacterium that has recently been found established in southern Europe, but whose potential impact and possible host range are still poorly documented in northern areas where the bacterium is not known to occur. To improve knowledge on the susceptibility of potential hosts of *X. fastidiosa* in northern Europe, a sentinel plantation of *Prunus domestica* cv. Opal, *Quercus petraea* and *Salix alba* was established in the *X. fastidiosa*-infected area of Majorca. In order to assess the circulation of the bacterium in the sentinel plot and around it, surveys of the local flora and insect vectors were carried out, as well as the planting of a network of rosemary “spy plants”. Symptomatic monitoring and molecular analyses were performed on the sentinel plants for four years. During these years, *X. fastidiosa* was never detected in our sentinel plants most likely because of the

low infectivity pressure recorded in the surroundings. This study underlines the complexity of conducting sentinel plantation assays combined with *X. fastidiosa* research, highlighting the need for long-term investigation and questioning the efficiency of the sentinel tool. However, this study is placed in perspective with other valuable sentinel plantations. It also highlights the complementarity of the tool and proposes elements to improve or re-orientate the implementation of future sentinel projects.

### Keywords

biological invasions, ex-patria planting, Majorca, northern Europe, pest risk analysis, *Prunus domestica*, *Quercus petraea*, *Salix alba*

## Introduction

The world sustainability is threatened by outbreaks of invasive pests and pathogens increasingly spreading around the globe (Simberloff et al. 2013; Diagne et al. 2021). These organisms largely travel to new areas through global trade, with living plants or with wood packaging material, which are considered as the main pathways of plant-related organism introductions (Kenis et al. 2007; Liebhold et al. 2012; Santini et al. 2013; Meurisse et al. 2019). These agents often expand by outcompeting native species because they are transported far from their natural enemies (“enemy release hypothesis”; Keane and Crawley 2002; Colautti et al. 2004), allowing them to allocate resources to growth and fecundity instead of defence, enhancing their fitness (“evolution of increased competitive ability” hypothesis; Blossey and Notzold 1995; Manfredini et al. 2013). They may trigger epidemics, sometimes on new hosts whilst they were less harmful to their native hosts, as they have not co-evolved with the new local plants that lack of specific defence mechanisms (Pimentel et al. 2001; Aukema et al. 2011). Apart from trade and globalisation, climate change and intensive land-use are also factors enhancing outbreaks by decreasing the resilience of the agricultural production systems and of forests (Walther et al. 2009; Bosso et al. 2016).

Preventing the introduction and the establishment of pests and pathogens in new areas is the most efficient tool for mitigating the consequences of a disease in terms of cost, biodiversity conservation and human impact (Barham et al. 2016). This includes the implementation of a pest risk analysis (PRA), which is an assessment giving biological, scientific and economic information on a particular organism (Aukema et al. 2011; Tomoshevich et al. 2013; EFSA PLH Panel 2018) to understand its potential impact and how it should be regulated (Parker et al. 1999; EC 2000; Liebhold et al. 2012). If considered harmful, the first measure taken to avoid its introduction might be its inclusion in a quarantine list implying either thorough inspections of imported plants before or after the importation, plant production in pest-free areas or sites of production or complete prohibition of trade or production of its native host plants (EC 2000).

However, these measures are not fully effective by themselves. Inspections can fail to intercept all the potential pests and pathogens travelling through plant trade (Kenis

et al. 2007; Eschen et al. 2015, 2017, 2019). First, these agents can be invisible to the naked eye because of their intrinsic nature or because they are in a latent form or in an endophytic stage on their traded hosts, leading to asymptomatic infections (Stergiopoulos and Gordon 2014; Migliorini et al. 2015). Secondly, despite the prioritisation of inspected organisms through PRA, the massive volume of traded materials makes the systematic control of each plant inoperable, with only batches being thoroughly examined (Britton et al. 2010; Eschen et al. 2015). Finally, PRA relies on prior awareness and knowledge of a pest and this knowledge is not always available; several agents, including non-catalogued taxa, harmless in their native region, are unknown to be invasive and pathogenic prior their introduction in a new land and escape controls (Brasier 2008; Britton et al. 2010; Tomoshevich et al. 2013; Vettrai et al. 2018). The few of them that manage to establish and cause significant damage are then often discovered too late to avoid outbreaks. Such is the case for some of the most damaging organisms of temperate forests that have occurred in recent years, which were unknown as pests prior to their introduction in a new area (Britton et al. 2010). Examples are the epidemics of Dutch elm disease caused by *Ophiostoma ulmi* Buisman and *O. novo-ulmi* Brasier that decimated billions of elm trees in Europe and America in the 20<sup>th</sup> century (Brasier and Buck 2001) or the massive damage to pines in Asia (Zhao et al. 2008) and Europe (Soliman et al. 2012) caused by the pine wood nematode, *Bursaphelenchus xylophilus* (Steiner & Bührer) Nickle, an organism well tolerated by its native pine hosts in North America (Akbulut and Stamps 2012).

A way to enhance knowledge about potentially damaging organisms to improve biosecurity systems would be to expose plants of interest out of their native range to study their susceptibility to local organisms in specific relevant locations, for example, a frequent plant exporting country (Roques et al. 2015; Barham et al. 2016). These plants would represent sentinels for their species in the foreign land. They provide an early warning for potential threats and additional information for PRA to set prevention measures and to know where the efforts for plant protection should be focused (Barham et al. 2016; Mansfield et al. 2019). An EPPO standard document was published in 2020, “PM 3/91 Standard on Sentinel woody plants” (EPPO 2020), to explain the approach and to provide guidance to carry out sentinel plant studies to identify new pest risks.

Sentinel plant research can be carried out by different ways (Britton et al. 2010). A first way is through botanical gardens and arboreta gathering a collection of specimens from all over the world, which are generally out of their area of origin and exposed to local agents. For such studies, the International Plant Sentinel Network (IPSN), working closely with National Plant Protection Organisations (NPPOs), was created. It connects the botanical gardens and arboreta staff around the world and gives them tools and expertise to monitor and to identify new pests and pathogens (Barham et al. 2016). Tomoshevich et al. (2013), for example, discovered 29 new pest-host associations (of which 18 were noticeably damaging for European trees) by studying European and Eurasian trees in Siberian gardens in Russia. However, in botanical gardens and arboreta, the number of representatives of each plant species is generally limited

(Roques et al. 2015), the trees are often large and hence difficult to examine in detail and they are usually subject to pesticide treatments or other management practices, which ensure plant health in the gardens (Eschen et al. 2019). Furthermore, gardens are often located in urban areas distant from the habitats of potential pests. All these reasons reduce the likelihood for an organism to reach and infect a specific plant species in an arboretum (Britton et al. 2010). A second way to conduct sentinel plant research is directly establishing actual plantations of exotic plants of interest in an environment where we want to study the impact of local pests and pathogens, the so-called “sentinel plantations” (Roques et al. 2015) or “ex-patria plantings” (Eschen et al. 2019). For example, Roques et al. (2015) and Vettraino et al. (2015) established two sentinel plantations of European tree species in China to investigate new pest-host associations potentially threatening to Europe that may emerge as a result of trade.

On the other hand, some well-known pathogens are still restricted to one part of the world and their potential host range in non-infected areas is uncertain and must be investigated. Such is the case of the phytopathogenic bacterium *Xylella fastidiosa* Wells et al., with more than 650 reported host plant species and for which the host range continues to extend as the bacterium enters new areas (EFSA 2022). While the threat of *X. fastidiosa* is definite for the European Mediterranean flora, the potential impact for northern areas is uncertain as most of the flora in these regions has never been exposed to the bacterium and probably contains many unreported hosts. The objective of our study was, therefore, to establish a sentinel plantation with European northern trees in a *X. fastidiosa* infected area in order to study the potential host range for these still-uninfected regions.

The gammaproteobacterium *X. fastidiosa* (Xanthomonadaceae) is strictly limited to the foregut of xylem sap-feeding insect vectors, mainly leafhoppers and spittlebugs (Hemiptera, Cicadomorpha) (Redak et al. 2004; Almeida et al. 2005; Chatterjee et al. 2008) and to the xylem vessels of its host plants. While many listed hosts are asymptomatic, the bacterium causes severe outbreaks on several crops, ornamental plants and shade trees generally provoking leaf-scorching that could lead to plant death (EFSA PLH Panel 2015). First limited to the Americas, the bacterium is currently regulated in Europe as a quarantine organism under the Council Directive 2000/29/EC (EC 2000). Between 2014 and today, the EuroPhyt database recorded 51 interceptions of *X. fastidiosa* in plants for planting and four interceptions of leafhoppers (EUROPHYT Online database 2022). Despite the border controls and EU prevention measures, a first focus of *X. fastidiosa* in Europe was discovered in 2013 in Apulian olive groves (Italy), for which more than 21 million olive trees were estimated to be affected in 2018 (Saponari et al. 2019b). The bacterium was then identified in mainland France, in Corsica (Denancé et al. 2017), in mainland Spain, in the Balearic Islands (Olmo et al. 2017), in another region of Italy (Tuscany) (Saponari et al. 2019a) and in Portugal (EUROPHYT Online database 2022). Divided into several subspecies (mainly subsp. *fastidiosa*, subsp. *multiplex* and subsp. *pauca*: Schaad et al. 2004) and more finely according to its sequence type (ST) (Scally et al. 2005; Yuan et al. 2010), 11 different STs were identified throughout Europe revealing multiple independent *X. fastidiosa*

introduction events (Cunty et al. 2022). Phylogeny studies allowed to date back the different entries of *X. fastidiosa* in the specific European regions, indicating entrance in the 1980s, 1990s and 2000s according to the area, i.e. well before the official identification of the pathogen's establishment on the continent. *Xylella fastidiosa* is, therefore, a perfect example of an organism escaping control due to the complexity of detection given the asymptomatic pool of hosts, the potentially long latent period limiting visual inspection and the number of reported and supposed/unreported hosts, as well as the lack of specific surveillance programmes and the limited availability of specific diagnostic tools in the past. Its movement into Europe has been caused in part by the trade of asymptomatic coffee plants imported from Latin America (EFSA PLH Panel 2015; Denancé et al. 2017).

However, it has been shown that eradication of *X. fastidiosa* may be complex if not impractical once it is well established and has reached a large geographical extent (Strona et al. 2017; EFSA PLH Panel 2019). Therefore, while entries can hardly be prevented, early detection is of prime importance to limit damage of outbreaks. The probability of early detection would increase by improving knowledge on where the bacterium is most likely to establish in order to conduct effective surveillance. In fact, performing detection tests on every plant in random areas is neither efficient nor conceivable as it would exceed any diagnostic capability considering the wide range of potential host plants (EFSA PLH Panel 2022). Targeting the main host plants and establishing a prioritisation list is essential to know where to focus resources and monitoring efforts.

To enhance knowledge on the susceptibility of potential hosts of *X. fastidiosa* in northern Europe, a sentinel plantation of northern plant species *Prunus domestica* cv. *Opal*, *Quercus petraea* and *Salix alba* was established in the *X. fastidiosa*-infected area of Majorca (Balearic Islands, Spain). There, the bacterium is considered widespread and well established. Three different STs belonging to two subspecies (*X. fastidiosa* subsp. *fastidiosa* ST1 and *X. fastidiosa* subsp. *multiplex* ST81 and ST7) have been identified on several hosts including wild olives, cultivated olives, almonds, grapes and figs (Olmo et al. 2021). They are mainly transmitted by the *Philaenus spumarius* Linnaeus (Aphrophoridae) vector and, to a lesser extent, by *Neophilaenus campestris* Fallén (Aphrophoridae) (López-Mercadal et al. 2021). This study experiments with the sentinel plantation tool in the case of *X. fastidiosa* research. The outcome questions the efficiency of the method, at least in this particular case and highlights the complexity of its implementation. However, it provides a methodology and several perspectives for future sentinel projects.

## Methods

### Preliminary tests and plant movement

The establishment and the monitoring of the sentinel plantation was achieved with the collaboration of the Applied Zoology and Animal Conservation group


of the University of the Balearic Islands (UIB). First, the agreement of the local government and the UIB authorities had to be obtained. Then, the plant material was bought at the Calle-Plant Nursery in Wetteren, Belgium. It consisted of dormant material: 30 *Salix alba* 0/1 80/120, 30 *Quercus petraea* 2/0 80/100 and 30 *Prunus domestica* cv. Opal 2 years grafted on Myrobolan or St Julien. Although all the plants were equipped with a phytosanitary certificate, *X. fastidiosa* specific detection tests were performed on several twigs of each plant to make sure the initial material was free of the bacterium. For this purpose, three branch parts of each plant were collected and bark peeled. They were chopped and their DNA was extracted according to the CTAB-based DNA extraction protocol specific for *X. fastidiosa* plant samples (“PM 7/24 (4) *Xylella fastidiosa*”, EPPO 2019). The detection was then performed by PCR (Minsavage et al. 1994). After this double check, the ninety plants were wrapped in hessian bags filled with wood chips and were brought by truck from Belgium to the UIB campus in Palma (Majorca, Balearic Islands) in March 2018. The chips were humidified during the 2-day trip to avoid root dryness.

### Location and establishment

The location of the plot was chosen with the UIB collaborators mainly based on the ease of connection to an irrigation system, as well as on the observation of *Philaenus spumarius* and *Neophilaenus campestris* nymphs on the ground vegetation and the presence of host plants, such as wild olive and almond trees. For the positioning of the plants in the plot, the JMP software was used to generate nine blocks, each one composed of three plants of each species randomly distributed (Fig. 1). The scheme was divided by blocks to take into account the potential gradients such as the slope, irrigation distribution or sunlight. The trees were planted directly into the ground to promote the growth of the root system and to enable them to survive throughout the season (Fig. 2). The soil was compact and rocky and was dug thanks to machines (Fig. 3). In every hole, about 20 litres of breeding soil were poured. The trees were separated from each other by 1.50 m and the whole plantation covered a total area of 144 m<sup>2</sup>. The irrigation system was established in the second year of the plantation. It consisted in three closed loops of pipe with one dripper per plant, allowing a constant pressure in all pipes and the same amount of water per plant (Fig. 1). The climatic data were followed through the season thanks to an HOBO device placed in the middle of the plantation.

### Exploring the surroundings

To monitor the circulation of the bacterium in the plot and around it, a 100-m demarcated area was organised around the plantation. In this area: i. a floristic inventory was carried out; ii. insect vectors were sampled; iii. a rosemary “spy plant” network was established (Fig. 4).




**Figure 1.** Scheme of the sentinel plantation of *Salix alba*, *Prunus domestica* cv. Opal and *Quercus petraea*. The dotted lines delimit nine blocks in which there are three plants of each species distributed randomly (JMP). The solid blue line is for the representation of the irrigation system consisting in three closed loops of pipe with one dripper per plant.



**Figure 2.** The sentinel plantation in Palma (Majorca, Balearic Islands) in May 2018.



**Figure 3.** Overview of the planting of the sentinel trees. Pictures highlighting the difficulty of establishing the sentinel plantation in the compact and rocky soil of the area.



**Figure 4.** Surroundings of the sentinel plantation **A** Google Earth view (Google Earth Pro, satellite image of 6 May 2021) of the UIB campus with the location of the sentinel plantation (purple square) and the 100-m demarcated area around the plantation (yellow circle) **B** scheme of the plantation and the demarcated area. In the demarcated area, a floristic inventory was carried out, insect vectors were sampled in the determined quadrat and a rosemary “spy plant” network was established by planting evenly seedlings around the plantation.

### Floristic inventory

To locate and assess the proportion of *X. fastidiosa* host plants in the area and to follow the eventual appearance of symptoms, a floristic inventory was carried out. It consisted

in identifying and mapping the tree layer of the demarcated area. An identification of the main herbaceous species was also performed with the help of local collaborators and of two determination keys for the local flora (Gil and Llorens 1999; Herbari Virtual del Mediterrani Occidental 2018).

### **Rosemary network**

A total of 44 *Rosmarinus officinalis* were planted around the campus: 32 plants evenly positioned in the demarcated area (Fig. 4B) and 12 plants in other places of the campus. The idea was to choose a robust plant adapted to local environmental conditions and which is quite susceptible to several subspecies of *X. fastidiosa*. Planting and regularly sampling these susceptible plants for bacterial detection provide a spy network allowing us to control the circulation of the bacteria in the vicinity of the plantation. The plants were bought in a local nursery in March 2018. They were first checked for *X. fastidiosa* presence with molecular tests before planting them, consisting of a CTAB-based DNA extraction followed by PCR of Minsavage et al. (1994). For sampling, about 15 leaves were collected on each plant, starting with symptomatic ones and were processed right away in the local laboratory. The mid-rib and the petiole were sectioned and the total DNA was extracted with the CTAB-based extraction procedure (EPPO 2019). The DNA samples were then sent to Belgium and were processed at UCLouvain by PCR of Minsavage et al. (1994) in the first three years and by real-time PCR of Harper et al. (2010) in the fourth year-final testing. In this final year, about five twigs per plant were collected as well and were processed in the same way.

### **Sowing ground vegetation**

As the planting of the sentinel plants with machines had removed the herbaceous layer in the sentinel plantation, which could prevent insects from reaching the trees, it was decided to re-sow grass in February of the second year to reconstitute this layer. The seed consisted of a universal mix of Asteraceae, Fabaceae and Poaceae.

### **Insect sampling and testing**

Insects were sampled with two objectives. On one hand, they were collected to be tested for *X. fastidiosa* presence by PCR (Minsavage et al. 1994) and quantitative PCR (Harper et al. 2010) to check for the circulation of the bacterium around the plantation. On the other hand, during the first year, the vector population density was assessed every month to determine the variability of the potential transmission during the season. For this study, the 100-m area around the plot was divided in 25 blocks (Fig. 4B). In each block, the same numbers of insect samples were taken. According to the development stage of the insect, the sampling method was adapted. For the nymphal stage, a 50 cm × 50 cm frame (0.25 m<sup>2</sup>) was thrown randomly four times in each block. The nymphs present in the surface delimited by the frame were counted.

In total, 84 samples were undertaken throughout the demarcated area and the number of nymphs/m<sup>2</sup> could be estimated. Although *X. fastidiosa* is lost after every moult, the nymphs can also become infected with it (Purcell and Finlay 1979; Redak et al. 2004). Therefore, in addition to the density study, three nymphs of *P. spumarius* and three nymphs of *N. campestris* were collected in each block for bacterial detection to determine if they already have an indication of the circulation of the bacteria in the plot. This quantity was chosen in order not to affect the vector abundance around the plot for the rest of the season.

Regarding insects at the adult stage, the sampling was carried out with sweeping nets. Two samples per block were taken in the ground layer, one sample corresponding to ten sweepings. The sweepings were undertaken homogeneously in each block in order to cover all the area. In total, 42 samples were taken throughout the demarcated area and the number of adult/sweep was measured. Again, only three insects per species (*P. spumarius* and *N. campestris*) were collected per block. Due to the small number of insects found in summer, the tree layer was also sampled. All the wild olive, almond and carob trees in the demarcated area were hit fifteen times with sweeping nets, distributed evenly on the plant in order to cover its entire attainable foliage surface. The number of adults per tree could be assessed.

The insects collected were placed at -20 °C, then stored in ethanol 70% and were sent to Belgium where they were processed. The eyes were removed and the DNA of the head together with the mouthparts was extracted using the CTAB-based protocol (EPPO 2019). The extracted DNA was then processed by PCR of Minsavage et al. (1994), by nested PCR of Cruaud et al. (2018) or by quantitative PCR of Harper et al. (2010).

## Sentinel plantation monitoring

Visual inspections were carried out for each sentinel tree. The appearance of *Xylella*-like symptoms was cautiously observed and wilting, shoot dieback, desiccation, defoliation or any change in leaf colour were reported. The evolution of the size of the different plants was also monitored, as well as the presence of *Xylella*-vectors or of other pests or organisms. In parallel, molecular analyses were performed on each plant. One sample per plant was collected, consisting of ten leaves per plant and 4–5 small twigs collected from all sides of the plant, but prioritising symptomatic areas if there were any. DNA extractions were carried out with the CTAB-based protocol (EPPO 2019) on leaf mid-ribs, on petioles and on the twigs after bark peeling and cutting them into small pieces. The DNA samples were then sent to Belgium where they were processed by PCR of Minsavage et al. (1994) in the first three years. In the final testing of the fourth year, two samples per plant were collected, one sample consisting of 10 different twigs distributed throughout the plant together with 10 to 20 leaves, always prioritising symptomatic parts. After extraction, they were processed by PCR of Minsavage et al. (1994), as well as by real-time PCR of Harper et al. (2010). No fertiliser was applied and no pruning was carried out in the winter, to allow the plants to develop naturally and not to cut potentially infected sections.

## Calendar of the establishment and monitoring of the plantation and demarcated area

The planning of the plantation monitoring during the four years is available in Table 1. The first year, it was decided to monitor the plantation and the demarcated area almost every month of the vector-season to assess the vector density fluctuation and to measure the rate of infection, if any, of the different plant species. In March, nymphs were sampled while, from May to October, insect adults were monitored. Several rosemary plants had desiccated already in May of the first year. Therefore, the dead ones were replaced in May and also in February of the following year. From the second year onwards, the sampling periods were chosen to correspond more or less to the beginning and the end of the highly infectious period of *X. fastidiosa* carried by the insect vectors, respectively, June and October, avoiding the aestivation periods of insects. The third year was impacted by the Covid-19 crisis and only one sampling campaign could be carried out in October 2020.


## Results

### Insight into surrounding plants

About 170 trees were inventoried: 134 carob trees, 31 wild olive trees, five almond trees and two pine trees. Their distribution can be observed in Fig. 5. The wild olive trees and the almond trees are both host plants of *X. fastidiosa*. Therefore, 36 host plants of the bacterium were identified in the 100 m around the plot (Fig. 5B). Amongst these host plants, 64% showed leaf scorching symptoms similar to those caused by *X. fastidiosa* (Fig. 5C). Concerning the ground vegetation, the identified plants were mainly: *Conium maculatum* (Apiaceae), *Foeniculum vulgare* (Apiaceae), *Cichorium intybus* (Asteraceae), *Dittrichia viscosa* (Asteraceae), *Galactites tomentosa* (Asteraceae), *Euphorbia medicaginea* (Euphorbiaceae) and many Poaceae (*Oryzopsis* sp. and others).

**Table 1.** Four-year schedule of the establishment and monitoring of the plantation and of the demarcated area.

Task	2018						2019		2020		2021	
	March	May	Jun	Jul	Sept	Oct	Feb	Jun	Oct	Oct	Jun	Oct
Sentinel establishment	■											
Rosemary network establishment	■	■					■					
Floristic inventory										C		
Vector density	■	■	■	■	■	■				O		
Vector sampling	■	■	■	■	■	■		■	■	V		
Sentinel plants monitoring and testing	■	■	■	■	■	■		■	■	I	■	■
Rosemary monitoring and testing	■	■	■	■	■	■		■	■	D	■	■
Sowing herbaceous vegetation							■					




**Figure 5.** Tree species inventory, host and health status around the sentinel plantation **A** map of the different tree species in the area of 100 m around the plantation. The pink square is the experimental plot (the sentinel plantation). The green dots are for *Olea europaea* var. *syliuestris* (wild olive tree), the blue dots for *Ceratonia siliqua* (carob tree), the pink dots for the *Pinus* sp. (pine tree) and the orange dots for the *Prunus dulcis* (almond tree) **B** map of the host status of the trees located in the area of 100 m around the plantation. The green dots are the non-host plants of *X. fastidiosa* and the yellow dots are the host plants of the bacterium **C** map of the symptomatic trees located in the area of 100 m around the plantation, presenting typical *X. fastidiosa* leaf scorches. The green dots are the asymptomatic plants, the yellow dots the symptomatic plants and the red dots the symptomatic plants that are host plants of the bacterium. The maps were created with the QGIS software with maps from Google Earth, Imagery 2018, DigitalGlobe.

Regarding the rosemary spy plants, molecular tests carried out over four years have not detected any bacteria in the collected samples. The rosemary plants suffered from the heat and many of them died. In May of the first year, the 12 rosemary planted in the campus were already all desiccated. The following year, they were replaced, as well as six rosemary plants located in the demarcated area. However, they did not last one year. Soil tilling performed in the demarcated area by the local gardeners also removed several plants from the ground. Only 12 out of 44 rosemary plants survived the four years of the experiment. In the first year, symptoms similar to those caused by *X. fastidiosa* already started to appear in May and, at the end of the first season, two thirds of the plants presented these symptoms, starting with chlorosis at the tip of the leaves, which extended to all the leaf surface and turned necrotic (Fig. 6).


### Insect sampling

Molecular tests carried out over four years have never detected any bacteria in the collected insects of the demarcated area.

During the first season, the amount of sampled insects of both species fluctuated depending on the month. This fluctuation can be observed in Fig. 7. In March, the foam produced by the nymphs could be easily observed and in total, 40 nymphs of *P. spumarius* (1.9 nymphs/m<sup>2</sup>, mainly at nymphal stage 3–4) and 89 nymphs of *N. campestris* (4.2 nymphs/m<sup>2</sup>, mainly at nymphal stage 2–3) were sampled (Fig. 7A, B).



**Figure 6.** Rosemary health state **A** sampled leaves of rosemary presenting *X. fastidiosa*-typical leaf scorch symptoms (May 2018) **B** dry and dead rosemary on the field (July 2018).




**Figure 7.** *Philaenus spumarius* (Ps) and *Neophilaenus campestris* (Nc) samples in 2018 in the 100 m area around the sentinel plantation **A** number of insects sampled through the different months. The striped pattern represents the nymphs and the plain pattern represents the adults **B** number of nymphs per m<sup>2</sup> sampled in March **C** number of adults per sweep sampled through the different months **D** number of adults per tree (wild olive, almond or carob tree) sampled during the different months.

The nymphs of *N. campestris* were always found on Poaceae, while those of *P. spumarius* were sampled on Asteraceae (*Carduus* sp.), Euphorbiaceae and other herbaceous plants. At the beginning of May, local collaborators observed nymphs of *P. spumarius* on one *S. alba* plant in the plantation, as well as two adults of *P. spumarius* on *P. domestica*.

At the end of May, the adult stage was already present and the sampling on the ground vegetation revealed less individuals than when nymphs were sampled the previous months. The number of adults per sweep was below one, with 0.04 *P. spumarius*/sweep and 0.03 *N. campestris*/sweep (Fig. 7C). In June, the herbaceous layer had dried and almost no insects were found in the ground vegetation. Very few insects were also sampled in the tree canopy. In September, more *P. spumarius* adults were sampled in the tree canopy (Fig. 7D); however, the number remained low with about 0.2 adults/tree. In October, new fresh herbs had grown and the highest number of *N. campestris* over the season was reached in the ground vegetation (0.13 adults/sweep), while a similar density as the one sampled in May was found for *P. spumarius* (0.03 adults/sweep).

The following years, the number of insects collected around the plantations varied between months and years (Fig. 8A, B) with a maximum in October 2020 of 0.06 *P. spumarius*/sweep and 0.13 *N. campestris*/sweep, sampled in the ground vegetation for both species. In total, four *P. spumarius* in October 2019, one *P. spumarius* in October 2020 and one *N. campestris* in October 2020 were found in the herbaceous layer of the sentinel plantation, showing that few insects were also circulating amongst the trees.




**Figure 8.** *Philaenus spumarius* (Ps) and *Neophilaenus campestris* (Nc) samples in 2019, 2020 and 2021 in the 100 m area around the sentinel plantation **A** the total number of collected adult insects **B** number of adults per sweep.

## The sentinel plants

Molecular tests carried out over four years have never detected any bacteria in the collected samples of the sentinel plants.

Nevertheless, first symptoms on *S. alba* already started to appear in June of the first year (2018) with some slight necrosis at the leaf margins of some of the plants. In July of that year, 78% (21/27 plants) of the willows had slight symptoms, while in October, 96% (26/27 plants) presented leaf necrosis starting from the tip, sometimes followed by chlorosis (Fig. 9A, B). Regarding *P. domestica*, slight chlorosis followed by necrosis at leaf margins started to appear in July 2018 on five of the plants (Fig. 9E). In October of the same year, ten plants had slight symptoms and two had moderate symptoms of chlorosis and necrosis of leaf margins. Finally, concerning *Q. petraea*, first typical necrosis on leaf margins started to appear in September of the first year. In October, these symptoms were more widespread affecting 30% of the plants (8/27 plants) and consisted of typical necrosis of leaf margins with a chlorotic halo (Fig. 9C), while two plants completely died. The following years, the same symptoms started to appear on the new growing leaves, mainly on *S. alba* and *Q. petraea*. On *P. domestica*, typical leaf symptoms were less frequent; however, this species presented




**Figure 9.** *Xylella*-like symptoms on the plants of the sentinel plantation in October of different years **A** on *Salix alba* in 2019 **B** on *Salix alba* in 2018 **C, D** on *Quercus petraea* in 2018 **E** on *Prunus domestica* in 2018.

more defoliation. The second year, the extremity of the principal stem of five plum trees and five willows started to die; for the three species, stem sprouts started to grow on 1–2 plants per species.

The summer of 2021 was declared the warmest recorded in Europe in the last 30 years, with severe heatwaves in the Mediterranean (Copernicus 2022). While the sentinel plants were already weakened by the last three hot summers despite the irrigation system, many of them died completely or partially this last year. Death was assigned after scraping the bark from several parts of the trunk. In total, 14 *S. alba* plants were completely dead and 13 had their main stem completely desiccated, but had developed sprouts at the bottom that were still living. The remaining leaves showed all symptoms of necrotic and chlorotic leaf margins. Three *Q. petraea* died and almost every remaining individual presented symptomatic leaves, while two of them had their main stem completely dead, but with living sprouts. Finally, two *P. domestica* died and about twenty of them had symptomatic leaves, which consisted of leaves turning red from the margins with a degraded colour, except for some leaves where the discoloured margins were quite delimited. About fifteen plants had between a quarter and a half of their main stem completely dead starting from the tip. Finally, the stem of two of them had completely died, leaving a second plant to grow from the variety Myrobolan, as the Opal variety was grafted on to this one. The size measured each year was not reported here because it was biased by the death, or partial death, of the main stem.

Concerning *Q. petraea*, damage caused by the herbivore *Lachnaia sexpunctata* Scopoli (Fig. 10A) in May–July 2018 forced us to put their foliage under a net (Fig. 10B) until mid-July to keep them alive, but this also resulted in their inaccessibility to *X. fastidiosa* insect vectors. A pesticide (Cypermethrin 10 ml/l) also had to be applied. The following years, the situation was better and the foliage could be exposed to the environment for all seasons. During the monitoring, fungal-like agents were also observed on the leaf surface of many individuals.



**Figure 10.** Herbivore damage on *Quercus petraea* **A** *Lachnaia sexpunctata* feeding on *Q. petraea* in the sentinel plantation **B** net on *Q. petraea* to avoid the herbivores eating their foliage.

## Discussion

During the four-year sampling and monitoring, *X. fastidiosa* was never detected in our sentinel plants nor in the collected insects. While it is rather positive not to have any infection of this quarantine pathogen, the duration of the plantation establishment did not allow us to answer the question of the potential host range of *X. fastidiosa*. In fact, besides the low infectivity pressure that had been observed on the plot, the absence of detected interaction between the bacterium and the sentinel plants does not mean that interaction could never occur (Mansfield et al. 2019) mainly given the highly specific conditions required by this plant pathogen. Instead, this study was an experimental work to learn how to combine sentinel plantation and research on *X. fastidiosa*, by exploring the constraints that were encountered to improve or redirect the method for future sentinel projects. In addition, the establishment of this plantation has provided valuable data on insect abundance and infection rates near the UIB campus and has enabled the implementation of other parallel experiments, while establishing a lasting international cooperation between the two universities.

### Complexity of sentinel plantations combined with *X. fastidiosa* research

Despite the publication of EPPO (2020) providing guidelines for sentinel studies, only two other assays that describe themselves as sentinel plantations have been reported in literature and both as part of the same project (Roques et al. 2015; Vettraino et al. 2015), while a third study can be characterised as one, even if it does not refer as such (Rathé et al. 2014). The sentinel plantations of Roques et al. (2015) and Vettraino et al. (2015) consisted of a four-year monitoring of five European tree species, including *Quercus* spp., which had been planted in China to investigate potential new host-pest/pathogen associations that could emerge in Europe through plant trade. While the experiments allowed the collection of valuable data and discovery of new associations, it already highlighted the complexity of the technique in terms of logistics and workload.

In our study, many constraints were faced and are reported in Table 2 with some perspectives on how the system could be improved to ease the implementation of the method. Our burdens started with permits and Italian administrations. In fact, the initial plan was to establish the plantation in the Apulian area where the first epidemic was declared. Ex-patria sentinel plantation studies require the movement and planting of non-native plants and they are, therefore, subjected to the host country's legislative and administrative procedures for importation and planting (EPPO 2020). After more than one year of back-and-forth e-mails to get approval from the Italian authorities, our request was transferred to our first correspondent. Therefore, the location of the plantation was changed to Majorca, where a good collaboration with UIB allowed us to obtain the agreement of the local authorities and the university, where the plantation was to be established, in about a month. A comparative view of the full procedural pathway between our first attempt in Apulia and Majorca can be viewed in Suppl. material 1. In addition, for administrative reasons, Roques et al. (2015) were unable to establish their plot in the initially optimal climatic zone where they wanted. In their

study, many plants were lost due to the delays in Chinese authorisations and imposed quarantine measures. Due to the common external border, plants with a European passport can circulate in Europe without restrictions and sentinel plantation *intra*-Europe should, therefore, be easier to implement (Vettrains et al. 2020). Furthermore, Vettrains et al. (2020) classified Europe as having low bureaucratic complexity concerning sentinel plantations compared to other non-European countries in a ranking they established according to the country's bureaucratic procedures. Surprisingly, Italy was considered the least complex European responding country, in contrast to what was experienced here. However, the current sensitive issue of *X. fastidiosa* in Italy has certainly not helped to speed up the procedures. On the other hand, the government of the Balearic Islands immediately accepted our request under certain conditions, which were the compliance with the norms in force in the territory regarding *X. fastidiosa* and the prohibition of planting *Polygala myrtifolia*, initially chosen as a spy plant for its high susceptibility to the bacterium. Vettrains et al. (2020) reported that most of the countries have restrictions on the import of certain plant species or genera, for example, Roques et al. (2015) were prohibited from planting *Pinus* spp. for their sentinel plantations in China. Finally, it is worth noting that we were not able to import plants collected in semi-natural environments, such as cuttings of *S. alba*, because of the difficulty of obtaining a phytosanitary passport for this type of material and all imported plants had to be purchased from Belgian nurseries in order to be certified.

The second challenge of this plantation was to keep the plants alive. The fact they were grown in an environment with different conditions including temperature and soil, brought different biotic and abiotic stresses. The life of these plants depended once again on the good cooperation on site. For example, the delay in the irrigation system establishment in the first year led the local staff to water the plants by hand every two days, carrying more than 80 litres of water in cans to the plantation. Furthermore, if they had not placed mesh covering the foliage for the herbivore *L. sexpunctata* that devoured the oak leaves, the plants would have died during the first year. However, despite constant monitoring by local collaborators, plant mortality increased from year to year and stress often led, especially in willows, to death of the main stem and the growth of new shoots at the bottom of the plant. This may have an impact on the outcome of the experiment, as the death of the potentially contaminated plant parts would lead to the death of the bacteria itself.

Here, the hurdles faced in sentinel plantation assays were coupled with the difficulties often encountered in *X. fastidiosa* studies. In fact, this bacterium is known to be fastidious for research including in its detection (Wells et al. 1987). Its concentration in plants and insects could be below the detection threshold of the different methods (Cruaud et al. 2018; EPPO 2019) and it is irregularly distributed in plants so may be missed during sampling, especially in asymptomatic plants (EFSA PLH Panel 2015; EPPO 2019). On the other hand, symptoms are not always reliable as they can easily be confused with symptoms triggered by other factors, such as drought (EFSA PLH Panel 2015). Therefore, it is more than likely that other causes, such as drought or soil stress, were responsible for the typical chlorosis and necrosis of the leaf margins

**Table 2.** Constraints and perspective of using sentinel plantation for *Xylella*-research. Constraints encountered in establishing a sentinel plantation in the case of a *Xylella fastidiosa* survey and perspectives for improving the implementation of the method.

Constraints	Perspectives
<b>Administrations</b>	
- Administrative procedures: Apulia vs. Majorca. Probably impacted by the sensitive issue of <i>X. fastidiosa</i> .	- Need for strong, organised and well-informed partnership. - Despite EPPO guidelines (2020), need for more homogenisation of admin. procedures and interpretation of the regulations at European level (and at global level through other intercontinental organisation/conventions), requiring to improve consideration towards sentinel plantations by increasing awareness of their usefulness.
<b>Legal framework of importing exotic plant material</b>	
- Complexity of obtaining a European passport for material collected in semi-natural environments.	- Need to simplify the procedures at national level for obtaining passports for scientific research purposes, under verification conditions of the plant material innocuity.
<b>Legal framework of <i>X. fastidiosa</i>, as a quarantine agent</b>	
- No movement of plant material from infected zones. - Removal of infected plants and vector control, decreasing infection pressure around the plantation.	- Need for a proper bio-molecular processing infrastructure on site. - Choosing a containment site and not an eradication site. - Need for further PRA exploration if special permits could be obtained for not uprooting infected flora for scientist research purpose or for maintaining plants under certain conditions, for example by placing an insect-proof net on the plants to prevent spread by vectors.
- Routing tests of local host plants not advised. - Restrictions of planting specific plant species ( <i>Polygala myrtifolia</i> ).	- Plantation of own susceptible spy plant network. - Or obtaining special permits for research purpose after PRA exploration. Need to verify plant innocuity and to sample them regularly to remove them as soon as possible in case of infection to prevent participating in the spread of the disease locally.
<b>Complexity to detect <i>X. fastidiosa</i></b>	
- Concentration below threshold of methods. - Irregular distribution in plants and asymptomatic plants. - Symptoms easily confused with ones due to other causes.	- Need to use several specific and sensitive detection methods (quantitative PCR, nested PCR...). - Multiply sampling from all sides of the plant (leaves and twigs). If symptoms detected, prioritising sampling of symptomatic parts. - Validation of bacterial presence only if detected with two different methods.
<b>Length of establishment</b>	
- Incubation period and length of establishment of <i>X. fastidiosa</i> . - Plants submitted to the unpredictability of natural conditions, with high dependence on abundance, host preference and prevalence of <i>X. fastidiosa</i> insect vectors.	- Long-term international financial and workload support. - Considering arboreta and botanical gardens studies. - Knowing the epidemiology of the exact sentinel location and choose a plot with high infective pressure (insect prevalence on site is measured on adults). - Considering targeted transmission experiments on sentinel plants with wild insects collected from naturally=infected areas.
<b>Pathosystem polymorphism</b>	
- Investigation only of potential pathosystems, based on local components. - For Majorca, strains: ST1, ST81, ST7; insect vectors: <i>P. spumarius</i> or <i>N. philaemus</i> ; local environmental conditions.	- Choosing location according to the strains one wants to test (extrapolating on current knowledge on which bacterial subspecies affect which plant genus can help, but it is not always accurate). - If no preferential subspecies, choosing location with the most strains present or multiply experiments to several areas. - Choosing location with the closest conditions to country of origin (environmental or insect vector population type).
<b>Abiotic and biotic stress for plants</b>	
- Other symptoms masking those of interest - Plant mortality limiting the experiment	- Irrigation system, eventual fertiliser application. - Fitting environmental conditions of native area if possible (in case of northern European countries more complicated with <i>X. fastidiosa</i> only occurring in southern Europe, thus, considering arboreta and botanical garden studies or targeted transmission experiments in infected areas).

observed on all three species in this study, especially for such plants used to colder temperature and more humid soil, even with the irrigation provided. While an undetectable low bacterial concentration can be questioned, several studies reported that high symptomatic responses were correlated with high bacterial loads (Holland et al. 2014; Saponari et al. 2017) suggesting a greater probability of detection if symptoms were due to *X. fastidiosa* infection.

Another parameter to consider when studying host susceptibility of *X. fastidiosa* is that the incubation period can be measured over months and years (EFSA PLH Panel 2019), indicating that time is a key element. For example, the survival time of Majorcan almond trees from bacterial infection to tree decline has been estimated around 14 years (Olmo et al. 2021). Sentinel plantation studies are already by themselves long-term assays and superimposing the potential time required for infection of the bacteria gives us an idea of how long it takes to conduct this type of experiment. However, the longer incubation period does not necessarily mean lower susceptibility to the bacterium itself, since many external factors can influence it, for example, the vector population. In fact, as *X. fastidiosa* is an insect vector-borne pathogen, its circulation and infection will depend upon the abundance, host preference and prevalence of its insect vectors, which are adding complexity to the system compared to other sentinel studies that would, for example, measure the direct impact of herbivores on leaves. Moreover, a particularity of diseases caused by *X. fastidiosa* is the polymorphism of the pathosystems. In fact, different strains and bacterial subspecies will act differently with the various xylem-feeding insect species and the different host species or cultivars, leading to very specific epidemics around the world (Pierce's disease, Citrus variegated chlorosis, Olive quick decline...) to almost no symptoms or to an endophytic presence. While the choice of the region in relation to the strains one wants to study is essential, this means that an absence or an endophytic interaction does not mean that other strains cannot be aggressive on the same plant species and cultivar. This means that there will only be an answer for a potential pathosystem related to the chosen region, but there are multitudes of other possibilities. The identified pathosystem will keep the adjective "potential" until the disease is not actually observed in the country of origin, as local environmental conditions or the presence of an effective vector will also have an impact.

A final element to be taken into account in the case of sentinel plantations with *X. fastidiosa* is the European regulation as a quarantine agent (Council Directive 2000/29: EC 2000) and the European containment and eradication measures imposed in case of detection (Commission Implementing Regulation (EU) 2020/1201: EC 2020) with the establishment of a demarcated area delimiting an infected zone of at least 50 m and a buffer zone varying in terms of kilometres depending on the situation. In the infected zone, eradication measures have to be undertaken consisting of the removal of all specified host plants of *X. fastidiosa*. However, in areas in which the bacterium is considered widely established including Apulia, Corsica and Balearic Islands, lighter containment measures may be implemented as eradication is no longer considered feasible. Nevertheless, these measures still imply the removal of all the infected

plants in the 50 m zone and an intensive surveillance within an area of at least 5 km radius together with vector control. These measures mean that, even in containment zones such as the Balearic Islands, the detection of an infected plant in the sentinel plantation would lead to the control of vector population in the area and to a decrease in the infection pressure around other plants of the plantation. Similarly, if the tested positive plant has to be removed immediately, the observation of symptom evolution and thus, the assessment of susceptibility is compromised, unless exceptional permits for scientific research are obtained. In this study, the problem did not arise because all plants tested negative. Nevertheless, we were still impacted by the consequences of the European legislation as, under the containment scenario in the Balearics, local government and UIB authorities did not advise systematic test of the host plants on the campus. In fact, a positive detection would have led to the uprooting of the campus vegetation, including, as mentioned before, our plantation if special permits were not issued. These measures are considered highly severe for an area where the bacterium is widespread and separated from other regions by the sea (Olmo et al. 2021). In areas infected by *X. fastidiosa*, the possibility of not having to remove infected plants in the field for scientific research purposes deserves further exploration in terms of PRA and bureaucratic procedure. Finally, for biosafety reasons related to quarantine organisms, plant samples cannot be moved and have to be processed on site, which again requires a good logistic, local collaboration and proper infrastructure.

### Necessity of knowing the epidemiology of the exact sentinel location


The implementation of a sentinel plantation when studying a specific pest or pathogen requires knowing well the epidemiology of the exact spot of the establishment, as local environmental components have a great impact on the outcome of the experiment (Kenis et al. 2018). The location chosen for this study was probably not optimal, as it was later evidenced that *X. fastidiosa* infection pressure was low and, thus, this certainly constitutes the main reason for the lack of positive detections in insects, spy and sentinel plants in the plot. When the plantation was established on the UIB campus, the prevalence and the epidemiology of the outbreak on the island were not yet well known, which is still the case in several regions where *X. fastidiosa* has recently been detected. Positive detections were reported on the campus about a hundred metres from the plantation on one *R. officinalis* plant and two olive trees (M. A. Miranda, personal communication) and the health state of host plants including declining almond trees, one of the main crop affected by *X. fastidiosa* on the Island, led us to suspect that the place was infected. However, due to the lack of systematic sampling after the declaration of the contention scenario in the Balearics, the presence of the bacterium could not be confirmed by testing. In addition, the quantity of nymphs sampled when choosing the location was 1.9 nymphs/m<sup>2</sup> for *P. spumarius* and 4.2 nymphs/m<sup>2</sup> for *N. campestris* in March, which is actually higher than the mean observed in the ground vegetation sampled through the Island. López-Mercadal et al. (2021) reported an average of about 0.22 nymphs/m<sup>2</sup> for *P. spumarius* in the peak of March and 0.005

nymphs/m<sup>2</sup> for *N. campestris* with differences between plots and years. In our plot, the prevalence of these nymphs was null. However, this information was not relevant as the infectivity is lost with each moult (Purcell and Finlay 1979) and prevalence, therefore, has to be measured on adult insects to have robust data.

After deepest outbreak investigations, it appeared that the east side of the Island towards Manacor was probably the most infected part, while the plantation was located to the west side of the Island. In fact, Gutiérrez Hernández and García (2018) mapped the positive records of *X. fastidiosa* detected in the Balearics by the Plant Health Section of the Department of Environment, Agriculture and Fisheries of the Government of the Balearic Islands and showed that most of the positive samples were concentrated on the east side with the highest densities in agricultural and residential areas close to the main communication routes. They stressed, however, that the conducted sampling strategy could have biased this distribution, for instance, because the samples could have been collected preferentially in these more accessible areas. Based on direct field observations and using Google street view, Moralejo et al. (2020) also mapped the distribution of *Xylella*-symptomatic almond orchards and their mortality across the Island, tracking their evolution since 2012 (Fig. 11). They showed a gradient from east to west, showing a moderate incidence on the site of the plantation. However, molecular testing of infected almond trees did not reveal a clear spatial pattern (Moralejo et al. 2020). In addition, highly variable incidence was encountered in different orchards (Olmo et al. 2021), hence the need of knowing the incidence and prevalence of vectors at the precise location of a sentinel plantation.

The density and prevalence of insect vectors are one of the drivers of *X. fastidiosa* infection and impact the temporal dynamics of symptom appearance (EFSA PLH Panel 2019), as multiple and independent infections could lead to an injection of a higher bacterial load and a decrease in the incubation period (Daugherty and Almeida 2009). The damage in the Balearics are the consequence of almost 20 years of infection (Moralejo et al. 2020), suggesting that the infection pressure could be too low to conduct sentinel plantation experiments. In fact, the abundance of nymphs and sampled adults, as well as the prevalence of insects are lower than the values encountered in the infected areas of Apulia where the outbreak was more drastic. A prevalence of 23% was reported in Majorca (López-Mercadal et al. 2021) compared to up to 71% detected in an Apulian olive grove (Cornara et al. 2016a). Similarly, higher densities of vectors were measured in Apulia with 7 to 39 nymphs of *P. spumarius*/m<sup>2</sup> in olive orchards (Bodino et al. 2019), about 7 adults/olive trees and 0.5 adults/sweep in weeds recorded during the respective seasonal peaks (Cornara et al. 2016b), however, with heterogeneity identified amongst the orchards studied (Bodino et al. 2019). In our plot, the adult density varied according to the seasonal estivation and ground drying pattern of Mediterranean regions (Cornara et al. 2016b; López-Mercadal et al. 2021). It barely reached a maximum of 0.04 *P. spumarius*/sweep in May and 0.2 *P. spumarius*/tree in September 2018, while the average reported through the Island was below 0.1 adults/sweep in ground cover, tree canopy and border vegetation (López-Mercadal et al. 2021). In addition, *N. campestris* was not considered as a significant vector due to its

Moralejo et al. (2020)



**Figure 11.** Almond leaf scorch incidence through Majorca. Map of the incidence of the almond leaf scorch disease and almond mortality within orchards across Majorca in 2012 and 2017 through field observation and Google view archives according to fig. 2 in Moralejo et al. (2020) (courtesy of E. Moralejo). The figure was adapted by adding the pink star at the localisation of the sentinel plantation.

very low presence on the tree canopy (López-Mercadal et al. 2021). Moreover, the soil around the plantation was ploughed almost every year, as common management on the Island, which, besides destroying several rosemary spy plants, probably decreased insect movement around the plot even with sowing of ground vegetation the second year. In fact, tillage is a technique of vector control reducing the number of vectors/m<sup>2</sup> (Bodino et al. 2019; EFSA PLH Panel 2019). In the study of Kenis et al. (2018), their plantation located at the edge of the forest took less time to be infested than another one situated in an agricultural-peri-urban area, highlighting again the impact of high local circulation of pests and pathogens on the time and outcome of the assay.

### Sentinel plantations as an efficient tool for *X. fastidiosa* research in specific situations

Even in locations with high infection pressure, the efficiency of the sentinel plantation in the case of *X. fastidiosa* host range investigation is questioned due to the ratio results/time-workload. Yet the sentinel plantation method is currently being used in Apulia for the screening of olive cultivars coming from various Mediterranean olive-growing

areas (Spain, Tunisia, Greece etc.) by exposing them to the natural pressure of inoculum in heavily-infected fields (XF-ACTORS 2017; Saponari et al. 2019b). The previous finding of the mild symptoms on the Leccino and FS17 olive cultivars adjacent to severely-affected orchards motivated the study (Boscia et al. 2017). Approximately 100 different genotypes were planted and are currently under evaluation in different plots, actually making the Apulian region home to one of the largest sentinel plantations of all time. This study is promising and is considered necessary for long-term management of *X. fastidiosa* in olive-growing regions as preliminary data show already differences in susceptibility in various cultivars (EFSA PLH Panel 2019; Saponari et al. 2019b). However, it highlights the long-term commitment required as the survey started in 2015 and is still ongoing. The project is part of a research programme funded by the European Union's Horizon 2020 Research and Innovation Programme, which explains how a project of this magnitude could be established and which underlines the need for long-term consistent international support for the implementation of such experiments. The success of this plantation, in addition to the selection of highly-infected plots, also comes from the fact that the tested potential hosts are related to the STs present in the environment. As the Apulian ST53 is highly aggressive on olives, it is obvious to carry out olive plant susceptibility in this area. However, other *X. fastidiosa* infected regions, such as Balearic Islands and Corsica, could be interesting to study the susceptibility to other STs, as three STs belonging to two subspecies co-exist in Majorca, while only one in the Apulian Region.

Thus, the Apulia study proved the usefulness of sentinel plantations in the context of *X. fastidiosa*. However, it would be less relevant to conduct these studies in certain situations. There should be, for example, similarities between the climatic conditions of the two regions involved in the sentinel studies to minimise the impact of external factors. So far, the bacterium has only been found established in southern Europe, in regions with a Mediterranean type of climate and these studies would, therefore, be less suitable for northern European countries, as differences in environmental conditions could lead to weakening or even death of the plants and to misidentification of the cause of potential symptoms. Nevertheless, this tool remains very valuable and should be considered for studies on *X. fastidiosa*, as other techniques for screening potential hosts of this pathogen are also discussed. Amongst these techniques, mechanical inoculation shows a low rate of success, even in susceptible hosts (Prado et al. 2008; EFSA PLH Panel 2019) as this method artificially reproduces infection while in the environment and only xylem-specialised insect vectors have the capacity to infect plants (Almeida et al. 2005). Working with insect vectors is, therefore, a more relevant way of conducting experiments. However, besides the biosafety risk it could represent for *Xylella*-free regions and the need for proper infrastructure, the very act of infecting an insect is a challenge. Other experiments consisting in grafting more than 400 olive genotypes on infected trees were conducted in parallel with the sentinel plantation in Apulia to short incubation period and time imposed by insect traits (Saponari et al. 2019b). However, in addition to also being an artificial way of infection, it requires the availability of appropriate infected graft material. Therefore, sentinel plantations have their advantages and have to be considered as valuable complementary tools in certain situations.

In these situations, this study has provided a complete methodology to monitor the bacterium circulation through the sentinel plants. The use of spy plants is certainly useful if sampling of susceptible vegetation is not possible in the nearby area. In other cases, sampling of local flora may be sufficient, although it does not ensure real-time circulation of the bacteria, as the current state of the local flora could be the result of infection from the past (Moralejo et al. 2020). The use of small perennial plants may facilitate sampling, as bacteria are distributed irregularly in the plant. The species or mix of species must be adapted to local conditions, susceptible to the bacterial strains being investigated and favoured by local vectors. In this study, *R. officinalis* was chosen as it was reported infected with the European STs of subsp. *multiplex* and subsp. *pauca* (ST6, ST7, ST53, ST80, ST81 and ST87) and was found infected in Majorca with the ST81 (EFSA 2022). In addition, in America, the bacterium was detected on this plant species close to *X. fastidiosa* subsp. *fastidiosa*-infected vines (Freitag 1951).

### Conducting sentinel studies differently to assess host range in northern countries

Sentinel studies can also be carried out differently to study host range in countries that cannot match closely the environmental conditions of the potential location. First, arboreta and botanical gardens are still an option for studying exotic host range in naturally-infected environments. However, as a detectable infection depends on the density and prevalence of *X. fastidiosa* insect vectors (Daugherty and Almeida 2009), the use of this method could also be discussed as these areas are often subjected to phytosanitary management. One advantage of these studies regarding *X. fastidiosa* would be that plants are grown in these sites for a long time, increasing the success concerning potential latent periods or low bacterial load potentially enabling detection. In addition, the study of Groenteman et al. (2015) has shown promising results for *X. fastidiosa* research by sampling in botanical gardens. They managed to discover 28 New Zealand plant species infected by *X. fastidiosa*, including several visited by the insect vector *Homalodisca vitripennis* Germar, in Californian botanical gardens where the disease is well established. They also found parasites capable of controlling the vector on these plant species with the aim of a biocontrol early-response strategy in case *H. vitripennis* invade New Zealand.

A second way would be to carry out transmission experiments in an insect-proof greenhouse with naturally-infected vectors in contaminated regions to bypass the problems of biosecurity imposed by *Xylella*-free areas and the difficulty of infecting insects. Compared to standard sentinel plantations, these experiments allow us to reduce the dependence on vector density and on insect feeding preferences. In fact, although *P. spumarius* is considered a polyphagous species and was observed feeding on the three studied sentinel plants in their area of distribution, it is possible that, in the sentinel country, these insects are more interested in native vegetation. Native plants could, therefore, compete with the exotic sentinel ones, potentially resulting in fewer vector feeding events decreasing the bacterial transmission probability. Even if vector preferences are biased and that natural conditions are, therefore, not fully met, these experiments can still be considered as sentinel studies since they consist in ex-patria plants

sent to study the impact of exotic organisms in areas in which they occur. This has been done in Majorca as a complementary experiment where 20 new cuttings of *S. alba* and of *P. tremula* have been sent from Belgium to the UIB campus. There, transmission experiments with naturally-infected *P. spumarius* were conducted in an insect-proof greenhouse and revealed positive infection on *S. alba*, proving the higher efficiency of the technique compared to sentinel plantation.

Finally, in-patria sentinel plantings (Eschen et al. 2019) or sentinel nurseries (sensu Vettraiño et al. 2017) consist of planting native traded plants without phytosanitary treatments on its own land to monitor pests and pathogens which could be spread through international trade (Vettraiño et al. 2017). They obviously do not have the same objective as ex-patria plantations that inform PRA of organisms that are not yet present in a given area. Rather, they consist of surveillance for a known pathogen for which possible entry and dispersal pathways have been identified (Mansfield et al. 2019) and they still represent valuable sentinel assays to be conducted with the aim of early detection of *X. fastidiosa* in new regions. The major difference with a standard commercial nursery is that no pest control measures are implemented on these plants (EPPPO 2020), so that it is possible for the vectors to reach the plants and for the plants to become infected if *X. fastidiosa* is introduced in the area. For this strategy to be effective, these plantations have to be established in strategic locations where the bacterium is the most likely to enter. The “plant for planting” pathway being the main entrance for exotic organisms including *X. fastidiosa* (Liebhold et al. 2012; EFSA PLH Panel 2018), their locations in/close to nurseries or other plant commercial places, would be relevant. In addition, these plantations must consist of known host plants that have a high probability to be the first infected when the bacterium enters an area and, if possible, to be highly susceptible for the infection to be visible and easily detectable. For example, the Auckland Botanic Garden has set up a sentinel plot of myrtle plants to detect the potential arrival of the myrtle rust (*Puccinia psidii* Winter) as early as possible in New Zealand, as the fungus was prevalent in Australia at the time (Barham et al. 2015). Similarly, one can imagine planting a network of *P. myrtifolia* near nurseries, previously tested for innocuity, which are regularly monitored for potential contamination by *X. fastidiosa*. Obviously, these susceptible plants should be tested carefully and regularly to provide the benefits of early detection while preventing them from serving as inoculum for disease establishment (Mansfield et al. 2019).

## Conclusion

In conclusion, this study is an experimental work highlighting that sentinel plantations are not easy to implement in the case of *X. fastidiosa*, but that they are complementary to other studies and that they could provide valuable information on host interactions when some conditions are met. This work proposes a methodology to monitor future sentinel plantations and it suggests other ways of conducting sentinel experiments for screening host range or for early detection of *X. fastidiosa* in new areas.

## Acknowledgements

We would like to acknowledge all the people who helped in the establishment of the sentinel plantation and the irrigation system and who contributed to maintain the plants alive, especially the UIB gardeners and technicians, Amélie Emond, Maria Antònia Tugores, Pau Mercadal, Carlos Barceló and Noelia Barros. We also want to thank all the people who irrigated manually with bravery the plantation during the first year. We are thankful to Sofia Delgado and Claudia Paredes for their laboratory support. Finally, we are grateful to UIB authorities and to the Government of the Balearic Islands for the authorisation to carry out these assays.

The research that yielded these results, was funded by the Belgian Federal Public Service of Health, Food Chain Safety and Environment through the contracts RF 19/6331 (XFAST project) and RT/7 XYLERIS 1 (XYLERIS project). NC was supported by the Foundation for Training in Industrial and Agricultural Research (FRIA, FNRS), and SH by the Belgian Federal Public Service of Health, Food Chain Safety and Environment.

## References

- Akbulut S, Stamps WT (2012) Insect vectors of the pinewood nematode: A review of the biology and ecology of *Monochamus* species. *Forest Pathology* 42(2): 89–99. <https://doi.org/10.1111/j.1439-0329.2011.00733.x>
- Almeida RPP, Blua MJ, Lopes JRS, Purcell AH (2005) Vector transmission of *Xylella fastidiosa*: Applying fundamental knowledge to generate disease management strategies. *Annals of the Entomological Society of America* 98(6): 775–786. [https://doi.org/10.1603/0013-8746\(2005\)098\[0775:VTOXFA\]2.0.CO;2](https://doi.org/10.1603/0013-8746(2005)098[0775:VTOXFA]2.0.CO;2)
- Aukema JE, Leung B, Kovacs K, Chivers C, Britton KO, Englin J, Frankel SJ, Haight RG, Holmes TP, Liebhold AM, McCullough DG, Von Holle B (2011) Economic impacts of non-native forest insects in the continental United States. *PLoS ONE* 6(9): e24587. <https://doi.org/10.1371/journal.pone.0024587>
- Barham E, Sharrock S, Lane C, Baker R (2015) An international plant sentinel network. *Sibbaldia: The International Journal of Botanic Garden Horticulture* (13): 83–98. <https://doi.org/10.24823/Sibbaldia.2015.75>
- Barham E, Sharrock S, Lane C, Baker R (2016) The International Plant Sentinel Network: A tool for regional and national plant protection organizations. *Bulletin OEPP/EPPO Bulletin. European and Mediterranean Plant Protection Organisation* 46(1): 156–162. <https://doi.org/10.1111/epp.12283>
- Blossey B, Notzold R (1995) Evolution of increased competitive ability in invasive nonindigenous plants: A hypothesis. *Journal of Ecology* 83(5): 887–889. <https://doi.org/10.2307/2261425>
- Bodino N, Cavalieri V, Dongiovanni C, Plazio E, Saladini MA, Volani S, Simonetto A, Fumara G, Di Carolo M, Porcelli, F, Gilioli G, Bosco D (2019) Phenology, seasonal abundance and stage-structure of spittlebug (Hemiptera: Aphrophoridae) populations in olive groves in Italy. *Scientific Reports* 9(1): 1–17. <https://doi.org/10.1038/s41598-019-54279-8>

- Boscia D, Altamura G, Ciniero A, Di Carolo M, Dongiovanni C, Fumarola G, Giampetruzzi A, Greco P, La Notte P, Loconsole G, Manni F, Melcarne G, Montilon V, Morelli M, Murrone N, Palmisano F, Pollastro P, Potere O, Roseti V, Saldarelli P, Saponari A, Saponari M, Savino V, Silletti MR, Specchia F, Susca L, Tauro D, Tavano D, Venerito P, Zicca S, Martel GP (2017) Resistenza a *Xylella fastidiosa* in diverse cultivar di olivo. L'Informatore Agrario 11: 59–63. <https://doi.org/10.5281/ZENODO.495708>
- Bosso L, Di Febbraro M, Cristinzio G, Zoina A, Russo D (2016) Shedding light on the effects of climate change on the potential distribution of *Xylella fastidiosa* in the Mediterranean basin. Biological Invasions 18(6): 1759–1768. <https://doi.org/10.1007/s10530-016-1118-1>
- Brasier CM (2008) The biosecurity threat to the UK and global environment from international trade in plants. Plant Pathology 57(5): 792–808. <https://doi.org/10.1111/j.1365-3059.2008.01886.x>
- Brasier CM, Buck KW (2001) Rapid evolutionary changes in a globally invading fungal pathogen (Dutch elm disease). Biological Invasions 3(3): 223–233. <https://doi.org/10.1023/A:1015248819864>
- Britton KO, White P, Kramer A, Hudler G (2010) A new approach to stopping the spread of invasive insects and pathogens: Early detection and rapid response via a global network of sentinel plantings. New Zealand Journal of Forestry Science 40: 109–114. <http://www.scopus.com/inward/record.url?eid=2-s2.0-77955440077&partnerID=tZOtx3y1>
- Chatterjee S, Almeida RPP, Lindow S (2008) Living in two worlds: The plant and insect lifestyles of *Xylella fastidiosa*. Annual Review of Phytopathology 46(1): 243–271. <https://doi.org/10.1146/annurev.phyto.45.062806.094342>
- Colautti RI, Ricciardi A, Grigorovich IA, MacIsaac HJ (2004) Is invasion success explained by the enemy release hypothesis? Ecology Letters 7(8): 721–733. <https://doi.org/10.1111/j.1461-0248.2004.00616.x>
- Copernicus (2022) Copernicus: Globally, the seven hottest years on record were the last seven; carbon dioxide and methane concentrations continue to rise. <https://climate.copernicus.eu/copernicus-globally-seven-hottest-years-record-were-last-seven> [Accessed on 10.06.2022]
- Cornara D, Cavalieri V, Dongiovanni C, Altamura G, Palmisano F, Bosco D, Porcelli F, Almeida RPP, Saponari M (2016a) Transmission of *Xylella fastidiosa* by naturally infected *Philaenus spumarius* (Hemiptera, Aphrophoridae) to different host plants. Journal of Applied Entomology 141(1–2): 80–87. <https://doi.org/10.1111/jen.12365>
- Cornara D, Saponari M, Zeilinger AR, de Stradis A, Boscia D, Loconsole G, Bosco D, Martelli GP, Almeida RPP, Porcelli F (2016b) Spittlebugs as vectors of *Xylella fastidiosa* in olive orchards in Italy. Journal of Pest Science 90(2): 521–530. <https://doi.org/10.1007/s10340-016-0793-0>
- Cruaud A, Gonzalez AA, Godefroid M, Nidelet S, Streito JC, Thuillier JM, Rossi JP, Santoni S, Rasplus JY (2018) Using insects to detect, monitor and predict the distribution of *Xylella fastidiosa*: A case study in Corsica. Scientific Reports 8(1): 1–13. <https://doi.org/10.1038/s41598-018-33957-z>
- Cunty A, Legendre B, de Jerphanion P, Dousset C, Forveille A, Paillard S, Olivier V (2022) Update of the *Xylella fastidiosa* outbreak in France: Two new variants detected and a new region affected. European Journal of Plant Pathology 163(2): 1–6. <https://doi.org/10.1007/s10658-022-02492-z>

- Daugherty MP, Almeida RPP (2009) Estimating *Xylella fastidiosa* transmission parameters: Decoupling sharpshooter number and feeding period. *Entomologia Experimentalis et Applicata* 132(1): 84–92. <https://doi.org/10.1111/j.1570-7458.2009.00868.x>
- Denancé N, Legendre B, Briand M, Olivier V, de Boissesson C, Poliakov F, Jacques MA (2017) Several subspecies and sequence types are associated with the emergence of *Xylella fastidiosa* in natural settings in France. *Plant Pathology* 66(7): 1054–1064. <https://doi.org/10.1111/ppa.12695>
- Diagne C, Leroy B, Vaissière AC, Gozlan RE, Roiz D, Jarić I, Salles JM, Bradshaw CJA, Courchamp F (2021) High and rising economic costs of biological invasions worldwide. *Nature* 592(7855): 571–576. <https://doi.org/10.1038/s41586-021-03405-6>
- EC [European Commission] (2000) Council Directive 2000/29/EC of 8 May 2000 on protective measures against the introduction into the Community of organisms harmful to plants or plant products and against their spread within the Community. *Official Journal of the European Communities* 169: 1–112. <http://data.europa.eu/eli/dir/2000/29/oj>
- EC [European Commission] (2020) Commission Implementing Regulation (EU) 2020/1201 of 14 August 2020 as regards measures to prevent the introduction into and the spread within the Union of *Xylella fastidiosa* (Wells et al.). *Official Journal of the European Union*. [http://data.europa.eu/eli/reg\\_impl/2020/1201/oj](http://data.europa.eu/eli/reg_impl/2020/1201/oj)
- EFSA PLH Panel [EFSA Panel on Plant Health] (2015) Scientific Opinion on the risks to plant health posed by *Xylella fastidiosa* in the EU territory, with the identification and evaluation of risk reduction options. *EFSA Journal* 13(1): 3989. <https://doi.org/10.2903/j.efsa.2015.3989>
- EFSA PLH Panel [EFSA Panel on Plant Health] (2018) Guidance on quantitative pest risk assessment. *EFSA Journal* 16(8): 5350. <https://doi.org/10.2903/j.efsa.2018.5350>
- EFSA PLH Panel [EFSA Panel on Plant Health] (2019) Update of the Scientific Opinion on the risks to plant health posed by *Xylella fastidiosa* in the EU territory. *EFSA Journal* 17(5): 5665. <https://doi.org/10.2903/j.efsa.2019.5665>
- EFSA (2022) Scientific Report on the update of the *Xylella* spp. host plant database – systematic literature search up to 31 December 2021. *EFSA Journal* 20(6): 7356. <https://doi.org/10.2903/j.efsa.2022.7356>
- EPPO [European and Mediterranean Plant Protection] (2020) PM 3/91(1) Sentinel woody plants: Concepts and application. *Bulletin OEPP. EPPO Bulletin. European and Mediterranean Plant Protection Organisation* 50(3): 429–436. <https://doi.org/10.1111/epp.12698>
- EPPO [European and Mediterranean Plant Protection Organization] (2019) PM 7/24 (4) *Xylella fastidiosa*. *Bulletin OEPP. EPPO Bulletin. European and Mediterranean Plant Protection Organisation* 49(2): 175–227. <https://doi.org/10.1111/epp.12575>
- Eschen R, Rigaux L, Sukovata L, Vettraino AM, Marzano M, Grégoire JC (2015) Phytosanitary inspection of woody plants for planting at European Union entry points: A practical enquiry. *Biological Invasions* 17(8): 2403–2413. <https://doi.org/10.1007/s10530-015-0883-6>
- Eschen R, Douma JC, Grégoire JC, Mayer F, Rigaux L, Potting RP (2017) A risk categorisation and analysis of the geographic and temporal dynamics of the European import of plants for planting. *Biological Invasions* 19(11): 3243–3257. <https://doi.org/10.1007/s10530-017-1465-6>

- Eschen R, O’Hanlon R, Santini A, Vannini A, Roques A, Kirichenko N, Kenis M (2019) Safeguarding global plant health: The rise of sentinels. *Journal of Pest Science* 92(1): 29–36. <https://doi.org/10.1007/s10340-018-1041-6>
- EUROPHYT Online database (2022) The European Union Notification System for Plant Health Interceptions. [https://ec.europa.eu/food/plant/plant\\_health\\_biosecurity/europhyt/interceptions\\_en](https://ec.europa.eu/food/plant/plant_health_biosecurity/europhyt/interceptions_en) [Accessed on 24.04.2022]
- Freitag JH (1951) Host range of the Pierce’s disease virus of Grapes as determined by insect transmission. *Phytopathology* 41(10): 920–934. <https://www.cabdirect.org/cabdirect/abstract/19521100495>
- Gil L, Llorens L (1999) Claus de determinacio de la flora balear. El Gall, Pollença, 1–293.
- Groenteman R, Forgie SA, Hoddle MS, Ward DF, Goeke DE, Anand N (2015) Assessing invasion threats: Novel insect-pathogen-natural enemy associations with native New Zealand plants in southern California. *Biological Invasions* 17(5): 1299–1305. <https://doi.org/10.1007/s10530-014-0804-0>
- Gutiérrez Hernández O, García LV (2018) Incidencia de *Xylella fastidiosa* en las Islas Baleares y distribución potencial en la península ibérica. *Investigaciones Geográficas [Esp]* (69): 55–72. <https://doi.org/10.14198/INGEO2018.69.04>
- Harper SJ, Ward LI, Clover GRG (2010) Development of LAMP and real-time PCR methods for the rapid detection of *Xylella fastidiosa* for quarantine and field applications. *Phytopathology* 100(12): 1282–1288. <https://doi.org/10.1094/PHYTO-06-10-0168>
- Herbari Virtual del Mediterrani Occidental (2018) Herbari Virtual del Mediterrani Occidental. [online] <http://herbarivirtual.uib.es/> [accessed repeatedly between March and June 2018]
- Holland RM, Christiano RSC, Gamliel-Atinsky E, Scherm H (2014) Distribution of *Xylella fastidiosa* in Blueberry Stem and Root Sections in Relation to Disease Severity in the Field. *Plant Disease* 98(4): 443–447. <https://doi.org/10.1094/PDIS-06-13-0680-RE>
- Keane RM, Crawley MJ (2002) Exotic plant invasions and the enemy release hypothesis. *Trends in Ecology & Evolution* 17(4): 164–170. [https://doi.org/10.1016/S0169-5347\(02\)02499-0](https://doi.org/10.1016/S0169-5347(02)02499-0)
- Kenis M, Rabitsch W, Auger-Rozenberg MA, Roques A (2007) How can alien species inventories and interception data help us prevent insect invasions? *Bulletin of Entomological Research* 97(5): 489–502. <https://doi.org/10.1017/S0007485307005184>
- Kenis M, Li H, Fan JT, Courtial B, Auger-Rozenberg MA, Yart A, Eschen R, Roques A (2018) Sentinel nurseries to assess the phytosanitary risks from insect pests on importations of live plants. *Scientific Reports* 8(1): 11217. <https://doi.org/10.1038/s41598-018-29551-y>
- Liebhold AM, Brockhoff EG, Garrett LJ, Parke JL, Britton KO (2012) Live plant imports: The major pathway for forest insect and pathogen invasions of the US. *Frontiers in Ecology and the Environment* 10(3): 135–143. <https://doi.org/10.1890/110198>
- López-Mercadal J, Delgado S, Mercadal P, Seguí G, Busquets A, Gomila M, Lester K, Kenyon DM, Pérez R, Paredes-Esquível C, Miranda MA (2021) Collection of data and information in Balearic Islands on biology of vectors and potential vectors of *Xylella fastidiosa* (GP/EFSA/ALPHA/017/01). EFSA Supporting Publications 18(10): 6925E. <https://doi.org/10.2903/sp.efsa.2021.EN-6925>
- Manfredini F, Grozinger CM, Beani L (2013) Examining the “evolution of increased competitive ability” hypothesis in response to parasites and pathogens in the invasive paper

- wasp *Polistes dominula*. *Naturwissenschaften* 100(3): 219–228. <https://doi.org/10.1007/s00114-013-1014-9>
- Mansfield S, McNeill MR, Aalders LT, Bell NL, Kean JM, Barratt BIP, Boyd-Wilson K, Teulon DAJ (2019) The value of sentinel plants for risk assessment and surveillance to support biosecurity. *NeoBiota* 48: 1–24. <https://doi.org/10.3897/neobiota.48.34205>
- Meurisse N, Rassati D, Hurley BP, Brockerhoff EG, Haack RA (2019) Common pathways by which non-native forest insects move internationally and domestically. *Journal of Pest Science* 92(1): 13–27. <https://doi.org/10.1007/s10340-018-0990-0>
- Migliorini D, Ghelardini L, Tondini E, Luchi N, Santini A (2015) The potential of symptomless potted plants for carrying invasive soilborne plant pathogens. *Diversity & Distributions* 21(10): 1218–1229. <https://doi.org/10.1111/ddi.12347>
- Minsavage GV, Thompson CM, Hopkins DL, Leite RMVBC, Stall RE (1994) Development of a polymerase chain reaction protocol for detection of *Xylella fastidiosa* in plant tissue. *Phytopathology* 84(5): 456–461. <https://doi.org/10.1094/Phyto-84-456>
- Moralejo E, Gomila M, Montesinos M, Borràs D, Pascual A, Nieto A, Adrover F, Gost PA, Seguí G, Busquets A, Jurado-Rivera JA, Quetglas B, García J de D, Beidas O, Juan A, Velasco-Amo MP, Landa BB, Olmo D (2020) Phylogenetic inference enables reconstruction of a long-overlooked outbreak of almond leaf scorch disease (*Xylella fastidiosa*) in Europe. *Communications Biology* 3(1): 1–13. <https://doi.org/10.1038/s42003-020-01284-7>
- Olmo D, Nieto A, Adrover F, Urbano A, Beidas O, Juan A, Marco-Noales E, López MM, Navarro I, Monterde A, Montes-Borrego M, Navas-Cortés JA, Landa BB (2017) First detection of *Xylella fastidiosa* infecting cherry (*Prunus avium*) and *Polygala myrtifolia* plants in Mallorca Island Spain. *Plant Disease* 101(10): 1820. <https://doi.org/10.1094/PDIS-04-17-0590-PDN>
- Olmo D, Nieto A, Borràs D, Montesinos M, Adrover F, Pascual A, Gost PA, Quetglas B, Urbano A, García J de D, Velasco-Amo MP, Olivares-García C, Beidas O, Juan A, Marco-Noales E, Gomila M, Rita J, Moralejo E, Landa BB (2021) Landscape Epidemiology of *Xylella fastidiosa* in the Balearic Islands. *Agronomy* 11(3): 473. <https://doi.org/10.3390/agronomy11030473>
- Parker IM, Simberloff D, Lonsdale WM, Goodell K, Wonham M, Kareiva PM, Williamson MH, Von Holle B, Moyle PB, Byers JE, Goldwasser L (1999) Impact: Toward a framework for understanding the ecological effects of invaders. *Biological Invasions* 1(1): 3–19. <https://doi.org/10.1023/A:1010034312781>
- Pimentel D, Mcnair S, Janecka J, Wightman J, Simmonds C, O'Connell C, Wong E, Russel L, Zern J, Aquino T, Tsomondo T (2001) Economic and environmental threats of alien plant, animal, and microbe invasions. *Agriculture, Ecosystems & Environment* 84(1): 1–20. [https://doi.org/10.1016/S0167-8809\(00\)00178-X](https://doi.org/10.1016/S0167-8809(00)00178-X)
- Prado SDS, Lopes JRS, Demétrio CGB, Borgatto AF, Almeida RPD (2008) Host colonization differences between citrus and coffee isolates of *Xylella fastidiosa* in reciprocal inoculation. *Scientia Agrícola* 65(3): 251–258. <https://doi.org/10.1590/S0103-90162008000300005>
- Purcell AH, Finlay A (1979) Evidence for Noncirculative Transmission of Pierce's Disease Bacterium by Sharpshooter Leafhoppers. *Phytopathology* 69(4): 393. <https://doi.org/10.1094/Phyto-69-393>

- Rathé AA, Pilkington LJ, Hoddle MS, Spohr LJ, Daugherty MP, Gurr GM (2014) Feeding and development of the glassy-winged sharpshooter, *Homalodisca vitripennis*, on Australian native plant species and implications for Australian biosecurity. PLoS ONE 9(3): e90410. <https://doi.org/10.1371/journal.pone.0090410>
- Redak RA, Purcell AH, Lopes JRS, Blua MJ, Mizell RF III, Andersen PC (2004) The biology of xylem fluid-feeding insect vectors of *Xylella fastidiosa* and their relation to disease epidemiology. Annual Review of Entomology 49(1): 243–270. <https://doi.org/10.1146/annurev.ento.49.061802.123403>
- Roques A, Fan JT, Courtial B, Zhang YZ, Yart A, Auger-Rozenberg MA, Denux O, Kenis M, Baker R, Sun JH (2015) Planting sentinel European trees in Eastern Asia as a novel method to identify potential insect pest invaders. PLoS ONE 10(5): e0120864. <https://doi.org/10.1371/journal.pone.0120864>
- Santini A, Ghelardini L, De Pace C, Desprez-Loustau ML, Capretti P, Chandelier A, Cech T, Chira D, Diamandis S, Gaitniekis T, Hantula J, Holdenrieder O, Jankovsky L, Jung T, Jurc D, Kirisits T, Kunca A, Lygis V, Malecka M, Marcais B, Schmitz S, Schumacher J, Solheim H, Solla A, Szabó I, Tsopelas P, Vannini A, Vettrains AM, Webber J, Woodward S, Stenlid J (2013) Biogeographical patterns and determinants of invasion by forest pathogens in Europe. The New Phytologist 197(1): 238–250. <https://doi.org/10.1111/j.1469-8137.2012.04364.x>
- Saponari M, Boscia D, Altamura G, Loconsole G, Zicca S, D’Attoma G, Morelli M, Palmisano F, Saponari A, Tavano D, Savino VN, Dongiovanni C, Martelli GP (2017) Isolation and pathogenicity of *Xylella fastidiosa* associated to the olive quick decline syndrome in southern Italy. Scientific Reports 7(1): 1–13. <https://doi.org/10.1038/s41598-017-17957-z>
- Saponari M, D’Attoma G, Abou Kubaa R, Loconsole G, Altamura G, Zicca S, Rizzo D, Boscia D (2019a) A new variant of *Xylella fastidiosa* subspecies *multiplex* detected in different host plants in the recently emerged outbreak in the region of Tuscany, Italy. European Journal of Plant Pathology 154(4): 1195–1200. <https://doi.org/10.1007/s10658-019-01736-9>
- Saponari M, Giampetruzzi A, Loconsole G, Boscia D, Saldarelli P (2019b) *Xylella fastidiosa* in olive in apulia: Where we stand. Phytopathology 109(2): 175–186. <https://doi.org/10.1094/PHTO-08-18-0319-FI>
- Sally M, Schuenzel EL, Stouthamer R, Nunney L (2005) Multilocus sequence type system for the plant pathogen *Xylella fastidiosa* and relative contributions of recombination and point mutation to clonal diversity. Applied and Environmental Microbiology 71(12): 8491–8499. <https://doi.org/10.1128/AEM.71.12.8491-8499.2005>
- Schaad NW, Postnikova E, Lacy G, Fatmi M, Chang CJ (2004) *Xylella fastidiosa* subspecies: *X. fastidiosa* subsp. *piercei*, subsp. nov., *X. fastidiosa* subsp. *multiplex* subsp. nov., and *X. fastidiosa* subsp. *pauca* subsp. nov. Systematic and Applied Microbiology 27(3): 290–300. <https://doi.org/10.1078/0723-2020-00263>
- Simberloff D, Martin JL, Genovesi P, Maris V, Wardle DA, Aronson J, Courchamp F, Galil B, García-Berthou E, Pascal M, Pyšek P, Sousa R, Tabacchi E, Vilà M (2013) Impacts of biological invasions: What’s what and the way forward. Trends in Ecology & Evolution 28(1): 58–66. <https://doi.org/10.1016/j.tree.2012.07.013>

- Soliman T, Mourits MCM, van der Werf W, Hengeveld GM, Robinet C, Lansink AGJMO (2012) Framework for modelling economic impacts of invasive species, applied to pine wood nematode in Europe. *PLoS ONE* 7(9): e45505. <https://doi.org/10.1371/journal.pone.0045505>
- Stergiopoulos L, Gordon TR (2014) Cryptic fungal infections: The hidden agenda of plant pathogens. *Frontiers in Plant Science* 5: 506. <https://doi.org/10.3389/fpls.2014.00506>
- Strona G, Carstens CJ, Beck PSA (2017) Network analysis reveals why *Xylella fastidiosa* will persist in Europe. *Scientific Reports* 7(1): 1–8. <https://doi.org/10.1038/s41598-017-00077-z>
- Tomoshevich M, Kirichenko N, Holmes K, Kenis M (2013) Foliar fungal pathogens of European woody plants in Siberia: An early warning of potential threats? *Forest Pathology* 43(5): 345–359. <https://doi.org/10.1111/efp.12036>
- Vettrano AM, Roques A, Yart A, Fan JT, Sun JH, Vannini A (2015) Sentinel trees as a tool to forecast invasions of alien plant pathogens. *PLoS ONE* 10(3): e0120571. <https://doi.org/10.1371/journal.pone.0120571>
- Vettrano AM, Li H-M, Eschen R, Morales-Rodriguez C, Vannini A (2017) The sentinel tree nursery as an early warning system for pathway risk assessment: Fungal pathogens associated with Chinese woody plants commonly shipped to Europe. *PLoS ONE* 12(11): e0188800. <https://doi.org/10.1371/journal.pone.0188800>
- Vettrano AM, Potting R, Raposo R (2018) EU Legislation on Forest Plant Health: An Overview with a Focus on *Fusarium circinatum*. *Forests* 9(9): 568. <https://doi.org/10.3390/f9090568>
- Vettrano AM, Santini A, Nikolov C, Grégoire JC, Tomov R, Orlinski A, Maaten T, Sverrisson H, Økland B, Eschen R (2020) A worldwide perspective of the legislation and regulations governing sentinel plants. *Biological Invasions* 22(2): 353–362. <https://doi.org/10.1007/s10530-019-02098-3>
- Walther GR, Roques A, Hulme PE, Sykes MT, Pyšek P, Kühn I, Zobel M, Bacher S, Botta-Dukát Z, Bugmann H, Czúcz B, Dauber J, Hickler T, Jarošík V, Kenis M, Klotz S, Minchin D, Moora M, Nentwig W, Ott J, Panov VE, Reineking B, Robinet C, Semchenko V, Solarz W, Thuiller W, Vilà M, Vohland K, Settele J (2009) Alien species in a warmer world: Risks and opportunities. *Trends in Ecology & Evolution* 24(12): 686–693. <https://doi.org/10.1016/j.tree.2009.06.008>
- Wells JM, Raju BC, Hung H-Y, Weisburg WG, Mandelco-Paul L, Brenner DJ (1987) *Xylella fastidiosa* gen. nov., sp. nov: Gram-negative, xylem-limited, fastidious plant bacteria related to *Xanthomonas* spp. *International Journal of Systematic Bacteriology* 37(2): 136–143. <https://doi.org/10.1099/00207713-37-2-136>
- XF-ACTORS (2017) Screening of olive cultivars for searching sources of resistance to *Xylella fastidiosa*. <https://www.xfactorsproject.eu/screening-cultivars-resistance-xf/> [Accessed on 06.06.2022]
- Yuan X, Morano L, Bromley R, Spring-Pearson S, Stouthamer R, Nunney L (2010) Multilocus sequence typing of *Xylella fastidiosa* causing Pierce's disease and oleander leaf scorch in the United States. *Phytopathology* 100(6): 601–611. <https://doi.org/10.1094/PHYTO-100-6-0601>
- Zhao BG, Futai K, Sutherland JR, Takeuchi Y (2008) Pine wilt disease, vol. 17. Springer, Tokyo, 459 pp. <https://doi.org/10.1007/978-4-431-75655-2>

## Supplementary material I

### **Administrative burdens and delays, a comparative view**

Authors: Noemi Casarin, Séverine Hasbroucq, Júlia López-Mercadal, Miguel Ángel Miranda, Claude Bragard, Jean-Claude Grégoire

Data type: PDF document

Explanation note: A comparative view of the full procedural pathway between the first attempt of implementing the sentinel plantation in Apulia and the second attempt in Majorca.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/neobiota.84.90024.suppl1>